What should I do?

If any of the early symptoms occur after eating shellfish, induce vomiting and get medical help immediately. There is no known antidote.

The Department of Health cannot guarantee the safety of recreationally harvested shellfish!

Shellfish can also be affected by other kinds of contamination. Be sure to gather shellfish away from pollution sources.

Questions?

Call your local health department or the Washington State Department of Health, Office of Food Safety & Shellfish Programs during office hours at (360)236-3330.



Marine Biotoxin Hotline is provided by the Washington State Department of Health Office of Food Safety & Shellfish Programs in cooperation with local health agencies.



TIDE

RED

Before you dig call

1-800-562-5632

or visit our web page at www.doh.wa.gov/ehp/sf/biotoxin.htm

Eating shellfish contaminated by a naturally occurring, toxic organism can cause serious illness, even death.

Each time you plan to gather shellfish from Puget Sound or other Washington beaches, call this 24-hour toll-free number or visit our web page for information on current biotoxin conditions.

PUBLIC HEALTH ALWAYS WORKING FOR A SAFER AND HEALTHIER WASHINGTON





How can I tell if shellfish are toxic?

The only foolproof way is through laboratory testing of the shellfish. Paralytic shellfish poison (PSP) and domoic acid poison toxins can be present with no apparent signs. Remember that boiling, cooking, or soaking the live shellfish in water does not remove or destroy the toxin.

What is the best way to keep informed on current conditions?

Call the Biotoxin Hotline
(1-800-562-5632) or access our
biotoxin web page at:
www.doh.wa.gov/ehp/sf/biotoxin.htm

Because conditions change rapidly, you should call or access our web site each time you plan to gather shellfish. These two resources provide information which can considerably reduce the risk of paralytic shellfish or domoic acid poisoning.

What about "red tide"?

Paralytic shellfish poison and domoic acid poison are rarely associated with a red tinge to the water. Don't assume the shellfish are safe if the water is NOT red. Reddish coloration of the water is more commonly associated with similar, but non-toxic, organisms. Both PSP and domoic acid can be present in large amounts when the water appears clear.

What causes the toxin?

When conditions such as sunlight, water temperature, and nutrients are favorable, a microscopic, poison-producing plankton can reproduce very rapidly in sea water. When shellfish feed on this plankton, they accumulate the toxin.

Are all shellfish potentially hazardous?

Clams, oysters, mussels, scallops, and moonsnails can become contaminated by the toxins. Crab may accumulate toxin in the gut. It is unknown if or how the toxin accumulates in finfish and other shellfish such as shrimp.

How will I know if I've eaten toxic shellfish?

The PSP toxin affects the nervous system. Early physical symptoms include tingling and numbness of the lips and tongue. Depending on how much toxin is consumed, symptoms may progress to tingling of the fingers and toes, difficulty breathing, and loss of control in the arms and legs. Death can occur if the respiratory system becomes paralyzed.

Domoic acid poisoning starts with gastrointestinal symptoms such as vomiting, nausea, diarrhea, and abdominal cramps within the first 24 hours. In more severe cases, neurological symptoms may develop within 48 hours that include headache, dizziness, confusion, and loss of short-term memory.

Domoic acid poisoning has been present above closure levels on coastal beaches in Washington State, but the Strait of Juan de Fuca and Puget Sound have not had toxin levels high enough to close those beaches.





Dear Colleague,

The Washington State Department of Health (DOH) is pleased to provide cameraready art for printing this educational material. To ensure that the original quality of the piece is maintained, please read and follow the instructions below and the specifications included for professional printing.

- Use the latest version. DOH materials are developed using the most current information available, are checked for clinical accuracy, and are field tested with the intended audience to ensure they are clear and readable. DOH programs make periodic revisions to educational materials, so please check this web site to be sure you have the latest version. DOH assumes no responsibility for the use of this material or for any errors or omissions.
- **Do not alter**. We are providing this artwork with the understanding that it will be printed without alterations and copies will be free to the public. Do not edit the text or use illustrations or photographs for other purposes without first contacting us. Please do not alter or remove the DOH logo, publication number or revision date. If you want to use a part of this publication for other purposes, contact the Office of Health Promotion first.
- For quality reproduction: Low resolution PDF files are intended for black and white or color desktop printers. They work best if you are making only one or two copies. High resolution PDF files are intended for reproducing large quantities and are set up for use by professional offset print shops. The high resolution files also include detailed printing specifications. Please match them as closely as possible and insist on the best possible quality for all reproductions.

If you have questions, contact:
Office of Health Promotion
P.O. Box 47833 Olympia, WA 98504-7833
(360) 236-3736

Sincerely, Health Education Resource Exchange Web Team

PRINTING SPECIFICATIONS

Title: Red Tide/Marine Biotoxin Hotline

Size: 3.667 x 8.5

Paper stock: 60# text Wausau Astrobrights Rocket Red

Ink color: Black

Special instructions: 2-sided printing.

DOH Pub #: 332-008